



LEADING
INNOVATION
IN ANIMAL HEALTH
FOR OVER 20 YEARS

Leading
INNOVATION
in animal health
for **over 20** years



SANFOVET


VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY




West Farm Company

Distributed by

WEST FARM COMPANY

 Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

 009647504660825 *** 009647507319748

 shorshvet@yahoo.com

 westfarm.info

Your
Success
is
Our
Priority





TABLE *of* **CONTENTS**

Liquid-based products

Powder-based products



West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Broad spectrum antibiotic

AZ.FLOTEC 25

COMPOSITION: Each 100ml contains

Florfenicol	25 g
Excipient q.s.f	100 ml

INDICATIONS:

Florfenicol has a broad-spectrum antibacterial activity that includes all organisms sensitive to florfenicol, against many gram-positive and gram-negative bacteria. It acts by inhibiting bacterial protein synthesis.

AZ.FLOTEC 25 is indicated for prevention and treatment of gastrointestinal and respiratory tract infections, caused by florfenicol sensitive micro-organisms such as Actinobaccillus spp., Pasteurella spp., Salmonella spp. and Streptococcus spp. in cattle and poultry.

ADMINISTRATION AND DOSAGE:

Dissolve into drinking water with the following concentration. Shake well before use.

Poultry: 1ml/25 kg of b.w or 1.000ml/5.000 litres (equivalent to 1 ml/5 litres) of drinking water for 3 - 5 days.

Medicated drinking water should be used within 24 hours. During treatment medicated drinking water should be the only source of drinking water.

WITHDRAWAL PERIOD:

Withdraw 21 days before slaughter in poultry.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Broad spectrum antibiotic

SANFO.QUINO 20% GOLD

COMPOSITION: Each 100ml contains

Enrofloxacin	20 g
Excipient q.s.f	100 ml

INDICATIONS:

Enrofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like Campylobacter, E. coli, Haemophilus,... which is used for the treatment of infections caused by bacteria sensitive to enrofloxacin such as respiratory diseases caused by Mycoplasma gallisepticum, Mycoplasma synoviae, intestinal and genital diseases caused by Escherichia coli in poultry as well as complication of infections in respiratory and GI tract of poultry: CRD, complex-CRD, colisepticaemia, salmonellosis, cholera, pasteurellosis, avian coryza, stress.

ADMINISTRATION AND DOSAGE:

Dissolve into drinking water with the following concentration.

Cattle, Sheep, Calf: 1ml/10kg b.w (20mg/kg b.w) of drinking water, for 3-5 consecutive days.

Poultry: 1.000ml/10 tons b.w (equivalent to 1ml/10kg b.w) of drinking water, for 3-5 consecutive days.

Small chick: 1000ml/4000 litres (equivalent to 1ml/4 liters) of drinking water, for 3-5 consecutive days.

Medicated drinking water should be used within 24 hours.

WITHDRAWAL PERIOD:

Withdraw 8 days before slaughter in poultry, cattle.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Timicosin 30%

AZ.TIMICOSIN ORAL

COMPOSITION: Each 100ml contains

Tilmicosin phosphate	30 g
Excipient q.s.f	100 ml

INDICATIONS:

AZ.TIMICOSIN ORAL has an antibacterial spectrum that is predominantly effective against Mycoplasma s.p., Pasteurella s.p. and Haemophilus s.p. and various Gram-positive organisms such as Corynebacterium spp.

AZ.TIMICOSIN ORAL is indicated for the control and treatment of respiratory infections associated with tilmicosin-susceptible micro-organism such as Mycoplasma spp., Pasteurella multocida, Haemophilus sp., Actinomyces pyogenes and Mannheimia haemolytica in livestock.

ADMINISTRATION AND DOSAGE:

Administration in drinking water or pet milk with the following recommended concentration:

Cattle: 1ml/25 kg b.w, administered twice per day, for 3 - 5 days.

Poultry, turkey: 300ml/1000 liters of drinking water, for 3 - 5 days.

Swine: 800 - 1600ml/1000 liters of drinking water, for 5 - 7 days.

Prepared solutions should be consumed within 24 hours.

To ensure correct dosage, concentration should be adjusted to the actual fluid intake.

If necessary, treatment can last for 10 - 21 days.

WITHDRAWAL PERIOD:

Withdraw 14 days before slaughter in cattle.

Withdraw 12 days before slaughter in poultry, turkey.

STORAGE:

Keep in dry, cool place and keep away from sunlight

Store in room temperature.

VETERINARY USE ONLY



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Broad spectrum antibiotic

ENROFLOXACIN LIQUID

COMPOSITION: Each 100ml contains

Enrofloxacin	10 g
Excipient q.s.f	100 ml

INDICATIONS:

Enrofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like Campylobacter, E. coli, Haemophilus,... which is used for the treatment of infections caused by bacteria sensitive to enrofloxacin such as respiratory diseases caused by Mycoplasma gallisepticum, Mycoplasma synoviae, intestinal and genital diseases caused by Escherichia coli in poultry as well as complication of infections in respiratory and GI tract of poultry: CRD, complex-CRD, colisepticaemia, salmonellosis, cholera, pasteurellosis, avian coryza, stress.

ADMINISTRATION AND DOSAGE:

Dissolve into drinking water with the following concentration

Poultry

- **Average dose:** 1ml/10 kg of b.w (10 mg/1 kg of b.w) or 1000ml/2000 litres (equivalent to 1ml per 2 liters) of drinking water, for 3 - 5 consecutive days.
- For the treatment of Colibacillosis or Salmonellosis for 5 - 7 consecutive days. If necessary, the treatment can be repeated after 5 days.

Calves, lambs, cattle

Average dose: 0,5ml/10 kg b.w for 3 days. Dose may be increased to 5 mg/kg for 5 days for salmonellosis and complicated respiratory disease. Medicated drinking water should be used within 24 hours.

WITHDRAWAL PERIOD:

Withdraw 8 days before slaughter in poultry, cattle.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,
Thuong Tin District, Hanoi, Vietnam
+84 969 68 0001
phongviavet@gmail.com
www.sanovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Natural herbal product improves breathing and respiratory system

MENTHOL PLUS

COMPOSITION: Each 100ml contains

Eucalyptus	4000 mg
Menthol	4000 mg
Thymol	2000 mg
Biotin	90 mg
Folic acid	20 mg
Methionine	3000 mg
Betaine	3000 mg
Excipient q.s.f	100 ml

INDICATIONS:

- Reduces the accumulation of mucus and the evacuation of respiratory cavities.
- Clears the passage of air into the lungs to improve breathing.
- Improves resistance to various respiratory infections.
- Promotes the growth rate and the consumption index.

ADMINISTRATION AND DOSAGE:

Dilute in drinking water to the following concentration.

Camel, cow, goat, sheep: 1-5ml/1liter of drinking water.

Poultry: 250ml/1000 liters of drinking water.

Spray method (applied to birds and poultry): dilute at a rate of 20ml/1 liter of water. Spray (fine nozzle) on the animal (aerial spray or aerosol generator) at a rate of 10 liters of solution to cover 2000 cubic meters.

WITHDRAWAL PERIOD:

Null.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Optimize hepatic function

HEPATOX FORTE

COMPOSITION:

 Each 100ml contains

Eucalyptus	4000 mg
Menthol	4000 mg
Thymol	2000 mg
Biotin	90 mg
Folic acid	20 mg
Methionine	3000 mg
Betaine	3000 mg
Excipient q.s.f	100 ml

INDICATIONS:

- Reduces the accumulation of mucus and the evacuation of respiratory cavities.
- Clears the passage of air into the lungs to improve breathing.
- Improves resistance to various respiratory infections.
- Promotes the growth rate and the consumption index.

ADMINISTRATION AND DOSAGE:

Dilute in drinking water to the following concentration.

Camel, cow, goat, sheep: 1-5ml/1liter of drinking water.

Poultry: 250ml/1000 liters of drinking water.

Spray method (applied to birds and poultry): dilute at a rate of 20ml/1 liter of water. Spray (fine nozzle) on the animal (aerial spray or aerosol generator) at a rate of 10 liters of solution to cover 2000 cubic meters.

WITHDRAWAL PERIOD:

Null.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Reduce mucus accumulation and clean the respiratory system

BROMREPSI

COMPOSITION: Each 100ml contains

Bromhexin HCl	2.000 mg
Menthol	800 mg
Excipient q.s.f	100 ml

INDICATIONS:

BROMREPSI increases serous incretion and clear mucus in respiratory tracts. As a result, mucus in the respiratory tracts are thinner to carry pathogens or any alien objects out of respiratory tracts.

BROMREPSI has antipyretic, analgesic and anti-inflammatory properties as well as:

- Acts as an expectorant to dissolve and loosen mucus (phlegm) in order to clear windpipe and ease breathing.
- Acts as a bronchodilator to widen airways to increase airflow in cases of severe viral diseases such as IB, ILT, Newcastle... or acute respiratory diseases such as ORT, ORD, CCRD....
- Prevents breathing problems caused by vaccination, adverse weather condition or transportation.

BROMREPSI does not inhibit other antibiotics' bioavailability when being combined. In fact, it helps enhance the concentration on lung tissue and increase the effectiveness of the treatment.

ADMINISTRATION AND DOSAGE:

Dilute in drinking water for 3 - 5 days consecutively with the following concentration: 1ml/4 litre of drinking water or 1 litre/4.000 litre of drinking water.


WITHDRAWAL PERIOD:

Before slaughter: 8 days.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY



SANFOVET

VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Solution for grow, fertility and health in animals

NUTRI OMEGA 3

COMPOSITION: Each 100ml contains

Omega 3 (fatty acids)	2.000 mg
Vitamin E	500 mg
Zinc Sulfate	300 mg
Sodium selenite	18 mg
Propylene Glycol	5.000 mg
Excipient q.s.f	100 ml

INDICATIONS:

Ruminants and other cattles

- Improves animals' health and well-being.
- Increases the milk production.
- Increases weight by fattening (calves-lambs).
- Improves meat quality.
- Acts as anti-inflammatory.
- Improves fertility, conception.
- Improves reproductive performance.
- Improves embryo survival rate.
- Enhance the calves' vitality.
- Shortens calving interval.
- Enhances immune system.

Poultry

- Increases egg production.
- Increases egg hatching rate.
- Enrich eggs with Ω 3 fatty acids.
- Improves the meat quality in broilers.
- Increases weight in broilers.
- Increases feed digestibility.

ADMINISTRATION AND DOSAGE:

Dilute in drinking water with the following recommended concentration.

Cattle-Camel: 15ml/head/day.

Sheep: 10ml/head/day.

Lamb: 5ml/head/day.

Poultry: 1.000ml/500 litres (equivalent to 2ml/1 litre) of drinking water.

WITHDRAWAL PERIOD:

Null.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,
Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Anticoccidial **AZ.DIAZURIL**

COMPOSITION: Each 100ml contains

Vitamin A	2.000.000 IU
Diclazuril	2.500 mg
Vitamin K3	500 mg
Excipient q.s.f	100 ml

INDICATIONS:

Diclazuril is a chemical anticoccidial, a benzeneacetonitrile derivative - synthetic compound of the triazinone family.

- **AZ.DIAZURIL** acts as an aid in the prevention of coccidiosis caused by Eimeria acervulina, E. brunetti, E. maxima, E. mitis, E. praecox, E. necatrix and E. tenella in broiler and layers.
- **AZ.DIAZURIL** acts an aid in the prevention of coccidiosis caused by Eimeria tenell, E. necatrix, E. acervulina, E. maxima, E.bruneti, E. mitis, E. mivati, E. hagani, E. praecox E. adenoides, E. meleagrimitis and E. gallopavonis in poultry.
- **AZ.DIAZURIL** also effective against coccidiosis caused by E. irresidua, E. media, E. flavescens, E. intestinalis, E. perforans, E. piriformis, E. stiedai, E. magna, E.exigua and E.coecicola in rabbits which stops the animal from gaining fat and improves reproductivity.

ADMINISTRATION AND DOSAGE:

Direct oral administration with the following recommended concentration: 1ml/25-30kg b.w/day (equivalent to 1000ml/25-30 tons b.w/day) or 2-3ml/10-15 litres of drinking water for 2-3 days.

WITHDRAWAL PERIOD:

Before slaughter: 5 days.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Ampicillin & Colistin

AMPI-COLI EXTRA

COMPOSITION: Each 100g contains

Ampicillin trihydrate	10 g
Colistin sulfate	25.000.000 IU
Excipient q.s.f	100 g

INDICATIONS:

For the treatment and prevention of bacterial infections in GI, respiratory and uro-genital tract such as colibacillosis, salmonellosis, white diarrhea (pullorum), fowl cholera and colisepticaemia.

AMPI-COLI Extra also treats secondary bacterial, infectious diseases during epidemic and relieve stress.

ADMINISTRATION AND DOSAGE:

Oral administration via drinking water or feed, for 3 - 5 days with the following recommended concentration:

- **Calves, lambs:** 1g/10 kg b.w of drinking water.
- **Poultry:** 1.000g/2.000 litres (equivalent to 1g/2 litres) of drinking water.

WITHDRAWAL PERIOD:

Withdraw 7 days before slaughter.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Gentamicin & Doxycycline SANFO.GENTADOX

INDICATIONS:

Doxycycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria like Bordetella, Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella, Staphylococcus and Streptococcus spp. Doxycycline is also active against Chlamydia, Mycoplasma and Rickettsia spp. The action of doxycycline is based on inhibition of bacterial protein synthesis. Doxycycline has good affinity towards the lungs and is therefore especially useful for treatment of bacterial respiratory infections.

Gentamicin sulphate belongs to the group of aminoglycosides and acts bactericidal against mainly Gram-negative bacteria like E. coli, Klebsiella, Pasteurella and Salmonella spp. The bactericidal action is based on inhibition of bacterial protein synthesis as well.

SANFO.GENTADOX can be used for the treatment of gastrointestinal and respiratory infections caused by micro-organisms sensitive to doxycycline and/or gentamicin in livestock.

ADMINISTRATION AND DOSAGE:

Dissolve into drinking water or mix with feed for 3 - 5 days.

- **Cattle:** 1g/10 kg of b.w.
- **Poultry:** 1.000g/1.000 litres (equivalent to 1g/1 litre) of drinking water or 1 g/5 kg of b.w.
- **Preventive dosage:** half the recommended concentration.

WITHDRAWAL PERIOD:

Withdraw 8 days before slaughter.
Withdraw 18 days before egg collection.
Withdraw 3 days before milk collection.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

COMPOSITION: Each 100g contains

Doxycycline hyclate	10 g
Gentamicin Sulfate	3 g
Excipient q.s.f	100 g



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

- Lien Phuong Industry Zone, Lien Phuong commune,
Thuong Tin District, Hanoi, Vietnam
- +84 969 68 0001
- phongviavet@gmail.com
- www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Tylosin & Gentamicin

AZ.GENTA-TYLOSIN

INDICATIONS:

For the treatment of gastrointestinal and respiratory infections caused by gentamicin and tylosin sensitive micro-organisms such as Bordetella, Campylobacter, Chlamydia, E. coli, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

ADMINISTRATION AND DOSAGE:

Oral administration via drinking water or feed, for 3 - 5 days with the following recommended concentration:

- **Cattle, goat:** 100g/500 kg of b.w/day or 1.000g/800 liters (equivalent with 10g/8 litres) of drinking water.
- **Poultry:** 100g/300 - 500kg of b.w/day or 1.000g/500 liters. (equivalent to 10g/5 litres) of drinking water.

WITHDRAWAL PERIOD:

Withdraw 7 days before slaughter.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

COMPOSITION: Each 100g contains

Tylosin tartrate	8 g
Gentamicin Sulfate	10 g
Vitamin A	500.000 IU
Excipient q.s.f	100 g



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Trimethoprim & Colistin SANFO.COLI-500

INDICATIONS:

For the treatment and prevention of infections of GI tract, or to suppress colonic flora such as colibacillosis, salmonellosis, pullorum, pasteurellosis in all animals and poultry, especially new-born animals.

ADMINISTRATION AND DOSAGE:

Dissolve into drinking water or mix with feed for 3 - 5 days with the following recommended dose:

- **Calves, goats and lambs:** 1g/15kg b.w (not more than 30 g per day per animal)
- **Poultry:** 1g/10kg b.w or 1.000g/2.000 litres (equivalent to 1g/2 litres) of drinking water.

WITHDRAWAL PERIOD:

Withdraw 5 days before slaughter.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

COMPOSITION: Each 100g contains

Trimethoprim	5 g
Colistin sulfate	120.000.000 IU
Excipient q.s.f	100 g



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Amoxicillin & Colistin

SANFO.AMOXY-COL 20%

INDICATIONS:

For the prevention and treatment of diseases caused by bacterial infection in livestock, including pneumonia, bacterial pneumonia caused by Hemophilus influenza, Colibacillosis, bacterial diarrhea, staphylococcosis, pasteurelosis, and salmonella, pneumonia, CRD, coryza, colibacillosis, staphylococcosis in animals.

ADMINISTRATION AND DOSAGE:

Oral administration via drinking water or feed, for 3 - 5 days with the following recommended concentration:

Poultry:

- Preventive dosage: 1.000g/8.000 litres (equivalent to 1g/8 liters) of drinking water or 100g/4.000kg of b.w daily.
- Curative dosage: 1.000g/4.000 litres (equivalent to 1g/4 liters) of drinking water or 100g/2.000kg of b.w daily.

WITHDRAWAL PERIOD:

Withdraw 5 days before slaughter.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

COMPOSITION: Each 100g contains

Amoxicillin trihydrate	20 g
Colistin sulfate	60.000.000 IU
Excipient q.s.f	100 g



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Resupplies electrolyte VIA-ELECTRAL

INDICATIONS:

- Resupplies electrolytes due to extensive loss of body fluid, such as diarrhea, enteritis, dehydration, persistent sweating, salivation, vomiting, bleeding,...
- Prevents retarded growth, impaired feed utilization, soft bones, muscle atrophy.
- Prevents weight loss during transportation.
- Restores electrolyte during stress condition, hot, high humidity, vaccination, regrouping.
- Stabilizes acid-base balance in the body in cases of acidosis, imbalance of K/Na ratio.

ADMINISTRATION AND DOSAGE:

Dissolve into drinking water or mix with feed for 3-5 days.

- **Poultry:** 1.000g/1.000 litres (equivalent to 1 g/1 liter) of drinking water.
- **Cattle, sheep, goats, horses:** 1.000g/100 litres (equivalent to 10g/1 liters) of drinking water.

WITHDRAWAL PERIOD:

Nil.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

COMPOSITION: Each 100g contains

Sodium hydrocarbonate	11.76 g
Sodium Chloride	15.2 g
Potassium Chloride	1.76 g
Excipient q.s.f	100 g



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

- Lien Phuong Industry Zone, Lien Phuong commune,
Thuong Tin District, Hanoi, Vietnam
- +84 969 68 0001
- phongviavet@gmail.com
- www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Feed supplement provide vitamins and trace elements

SUPER ES20

INDICATIONS:

Super ES 20 is considered a dietary supplement to compensate for the deficiency of vitamin E and selenium, as well as zinc.

ADMINISTRATION AND DOSAGE:

Mix in drinking water for 3-5 days.

- **Camel, cow:** 2.5 grams/liter/day/head.
- **Sheep, goat:** 1.5 grams/liter/day/head.
- **Poultry:** 1.000g/10.000 litres. (equivalent to 1g/10 litres) of drinking water.

WITHDRAWAL PERIOD:

Null.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

COMPOSITION: Each 100g contains

Vitamin E	20 g
Selenium (Sodium selenite)	50 mg
Zinc (Zinc Sulfate)	10 g
Excipient q.s.f	100 g



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,
Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Providing energy and accelerating recovery

KC GLUCOVIT

INDICATIONS:

KC GLUCOVIT is used for sick cattle and poultry that are experiencing the following symptoms: IBD (Infectious bursal disease), CRD (Chronic respiratory disease), Gumboro disease, stress,...

KC GLUCOVIT helps building immune system to fight infectious diseases and relieve stress during environmental changes and vaccination.

KC GLUCOVIT also helps preventing and treating inflammation, haemorrhagic diathesis, exudative, allergy and intoxication.

KC GLUCOVIT improves fertility, hatchability and eggshell quality.

ADMINISTRATION AND DOSAGE:

Oral administration via drinking water with the following recommended dosage: 1.000g/500 litres (equivalent to 10g/5 liters) of drinking water, for 5 - 7 days.

WITHDRAWAL PERIOD:

Nil.


STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

COMPOSITION: Each 100g contains

Glucose	98 g
Vitamin K3	40 mg
Vitamin C	1200 mg
Excipient q.s.f	100 g



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,
Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Beneficial bacteria, vitamins and electrolyte powder

AZ.BIOZYME ONE

INDICATIONS:

AZ.BIOZYME ONE is a food supplement in the form of a soluble powder containing bacteria and vitamins that help.

- Creates the optimal environment for the growth and activity of beneficial bacteria in the intestine.
- **AZ.BIOZYME ONE** helps improve the performance of the digestive system resulting from the use of antibiotics.
- Improving the smell in the barn.

ADMINISTRATION AND DOSAGE:

Dissolve in drinking water or mix with feed for 3 - 5 days.

- Camel, cow, goat, sheep: 5g/1 liter/day/head or 0.5 kg/1 ton of feed.
- Poultry: 1.000g/1.000 litres (equivalent to 1 g/1 liter) of drinking water or 0.5 kg/1 ton of feed.

WITHDRAWAL PERIOD:

Null.


STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

COMPOSITION: Each 100g contains

Lactobacillus acidophilus	1.000.000 CFU
Lactobacillus reuteri	1.000.000 CFU
Lactobacillus casei	1.000.000 CFU
Lactobacillus rhamnosus	1.000.000 CFU
Bacillus subtilis	1.000.000 CFU
Protease	33.000 UI
Vitamin C	1.000 mg
Sodium (Na)	2.450 mg
Potassium (K)	1.550 mg
Excipient q.s.f	100 g



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Oral nutritious supplement powder **AMINOVIT + MINERAL**

COMPOSITION: Each 1.000g contains

Vitamin A	6.700.000 IU	Calcium (Ca)	440 mg
Vitamin D3	53.000 IU	Iron (Fe)	8.000 mg
Vitamin E	12.000 mg	Copper (Cu)	825 mg
Vitamin B1	440 mg	Manganese (Mn)	4.000 mg
Vitamin B2	1.600 mg	Zinc (Zn)	9.000 mg
Vitamin B6	950 mg	Iodine (I2)	60 mg
Vitamin B9	1.500 mg	Cobalt (Co)	10 mg
Vitamin B12	1.100 mg	Selenium	10 mg
Vitamin C	5.400 mg	L-Lysine	8.000 mg
Vitamin PP	5.400 mg	Chlorine	7.500 mg
Vitamin K3	590 mg	Excipient q.s.f	1.000 g

INDICATIONS:

AMINOVIT+MINERAL is a soluble powder supplement containing vitamins, amino acids and minerals which helps:

- Creating the optimal environment for beneficial bacteria in the intestine during growth for animal.
- Stimulates appetite, help them in gaining weight fast, stimulate rapid growth in mass and muscle which would provide high quality meat, also saving cost by shorten feed conversion ratio (FCR).
- Improves immune system in animal, when vaccines is applied, the effect will intertwine for a better result.
- Helps animals to gain good physique, improve semen quality and reproductive efficiency.

WITHDRAWAL PERIOD:

Null.

STORAGE:


Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

ADMINISTRATION AND DOSAGE:

Dissolve into drinking water or mix with feed for 3-5 days with the following recommended dosage:

- **Camel:** 30 g/head/day.
- **Cows, sheep, goats:** 10-30g/head/day.
- **Poultry:** 1.000-2.000g/1.000 litres (equivalent to 1 - 2 g/1 litre) of drinking water or 1.000g/0.5 tons of feed.



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Assure hatching rate, quality of embryos, and eggs

SUPER.EGG

COMPOSITION:

Each 1.000g contains

Oxytetracycline (HCL)	60.000 mg	Calcium Gluconat	75.000 mg
Vitamin A	4.000.000 IU	Magnesium sulfate	11.200 mg
Vitamin D3	700.000 IU	Zinc Sulfate	11.100 mg
Vitamin E	1.000 mg	Sodium Chloride	15.000 mg
Vitamin B2	2.500 mg	DL-Methionine	25.000 mg
Vitamin B12	5.000 mcg	L-Lysine	45.000 mg
Calci D-pantothenate	6.000 mg	Vitamin K3	1.500 mg
Vitamin PP	15.000 mg	Excipient q.s.f	1.000 g

INDICATIONS:

- Super.Egg is an exclusive product for layers which helps to prolong rearing period.
- Super.Egg ensures that eggs and embryo quality as well as hatching rate is at high level.
- Super.Egg also helps preventing stress during transportation and environmental changes.

WITHDRAWAL PERIOD:

Nil.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

ADMINISTRATION AND DOSAGE:

Dissolve into drinking water or mix with feed for 3-5 days with the following recommended dosage: 1.000g/1.000 litres (equivalent to 1g/1 litre) of drinking water or 100 g/500 kg of b.w per day.
When the flock is suffering from stress or sickness, administer 1.000g/300 litres (equivalent to 10g/3 litres) of drinking water or 100g/300 kg of b.w per day for 5 - 7 days consecutively.



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Stimulates and prolong high yield of egg production

VITASTRONG

COMPOSITION: Each 1.000g contains

L-Lysine	25 g	Vitamin K3	10 g
Potassium iodide	100 mg	Vitamin C	50 g
Copper Sulfate	1500 mg	Vitamin B1	10 g
Cobalt Sulfate	100 mg	Vitamin B2	10 g
Iron Sulfate	5000 mg	Nicotinic Acid	75 g
Manganese Sulfate	10 g	Ca-d-pantothenate	25 g
Zinc Sulfate	10 g	Vitamin B6	10 g
D,L- Methionine	10 g	Vitamin B12	25.000 mcg
Vitamin A	25.000.000 IU	Folic Acid (B9)	200 mg
Vitamin D3	3.000.000 IU	Excipient q.s.f	1.000 g
Vitamin E	50 g		

INDICATIONS:

- Stimulates and prolongs egg production. Increases blastocyst rate to maximize hatchability rate.
- Helps balance livestock's diet where nutrient deficiencies exist and fasten their recovery.
- Supplies minerals and amino acids to regulate digestive function so that livestock can remain their vigorous appetite.
- Reduces stress due to vaccination and aids in the treatment of diseases caused by environmental changes.

WITHDRAWAL PERIOD:

Nil.

STORAGE:


Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

ADMINISTRATION AND DOSAGE:

Oral administration via drinking water or feed with the following concentration: 100 g/500 kg b.w per day or 1 kg/1000 litres of drinking water every day during rearing stage.

In case of stress, sudden drop in egg production/quality and/or low blastocyst rate: double the recommended dosage for 3-5 days consecutively.



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,

Thuong Tin District, Hanoi, Vietnam

+84 969 68 0001

phongviavet@gmail.com

www.sanfovet.com





West Farm Company

Distributed by

WEST FARM COMPANY

Makhmur road
60 Meter st Near security center
ERBIL-IRAQ

009647504660825 *** 009647507319748

shorshvet@yahoo.com

westfarm.info

Instantly reduce fever and fasten recovery

PARAMAX C

COMPOSITION: Each 100g contains

Paracetamol	10 g	Excipient q.s.f	100 g
Vitamin C	5 g		

INDICATIONS:

- Relieve pain and reduce fever, increase resistance against bacterial and viral diseases.
- It contains vitamin C which plays important role in maintaining the activity and vitality of the animals' body, and increase the resistance especially after stress and fatigue.
- It is used for stress reaction caused by vaccinations in poultry.
- It can be combined during treatment with antibiotics.

WITHDRAWAL PERIOD:

Nil.

STORAGE:

Keep in dry, cool place and keep away from sunlight
Store in room temperature.

VETERINARY USE ONLY

ADMINISTRATION AND DOSAGE:

Oral administration via drinking water or feed with the following recommended concentration:

- **Poultry:** 1.000g/1.000 litres (equivalent to 1g/1 litre) of drinking water.
- **Cattle:** 1g/8-10 kg b.w/day.



VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

Lien Phuong Industry Zone, Lien Phuong commune,
Thuong Tin District, Hanoi, Vietnam
+84 969 68 0001
phongviavet@gmail.com
www.sanfovet.com






THE **BRAND**
FOR **QUALITY**
AND **PRESTIGE**




SANFOVET


Contact Us

VIET ANH JOINT VENTURE INVESTMENT JOINT STOCK COMPANY

 Lien Phuong Industry Zone, Lien Phuong commune,
Thuong Tin District, Hanoi, Vietnam

 +84 969 68 0001

 phongviavet@gmail.com

 www.sanfovet.com

